

Quantitative results for Bruck iterations of demicontinuous pseudocontractions

Daniel Körnlein*

Department of Mathematics
Technische Universität Darmstadt
Schlossgartenstraße 7, 64289 Darmstadt, Germany

October 4, 2016

Abstract

Our first result is a rate of metastability in the sense of Tao for Bruck's iteration scheme for demicontinuous pseudocontractions in Hilbert space, extracted from Bruck's original proof. This result generalizes earlier work in the ongoing program of proof mining from Lipschitzian to demicontinuous pseudocontractions. Our second main result is a metastable version of asymptotic regularity under the additional assumption that the underlying operator is norm-to-norm uniformly continuous on bounded subsets. These results (and their intermediate versions given in this paper) provide a thorough quantitative analysis of Bruck's iteration scheme for pseudocontractions in Hilbert space.

1 Introduction

Let X be a normed linear space and $S \subseteq X$ be a subset of X . In 1967, Browder introduced an important generalization of the class of nonexpansive mappings, namely the *pseudocontractive* mappings $T : S \rightarrow S$ defined by

$$\forall u, v \in S \forall \lambda > 1 ((\lambda - 1)\|u - v\| \leq \|(\lambda I - T)(u) - (\lambda I - T)(v)\|),$$

where I denotes the identity mapping.

Apart from being a generalization of nonexpansive mappings, the pseudocontractive mappings are also closely related to accretive operators, where an operator A is called accretive if for every $u, v \in S$ and for all $s > 0$,

$$\|u - v\| \leq \|u - v + s(Au - Av)\|.$$

Observe that T is pseudocontractive if and only if $I - T$ is accretive. Therefore, any fixed point of T is a root of the accretive operator $I - T$.

In a Hilbert space, T is pseudocontractive if and only if

$$\forall u, v \in S (\langle Tu - Tv, u - v \rangle \leq \|u - v\|^2)$$

(see e.g. [2]).

In [1], Bruck introduced the following iteration schema for pseudocontractive mappings:

Definition 1.1 ([1]). Let C be a nonempty convex subset of a real normed space and let $T : C \rightarrow C$ be a pseudocontraction. Let $(\lambda_n), (\theta_n)$ be sequences in $[0, 1]$ with $\lambda_n(1 + \theta_n) \leq 1$ for all $n \in \mathbb{N}$. The **Bruck iteration scheme** with starting point $x_1 \in C$ is defined as

$$x_{n+1} = (1 - \lambda_n)x_n + \lambda_n T x_n - \lambda_n \theta_n (x_n - x_1).$$

*The author has been supported by the German Science Foundation (DFG Project KO 1737/5-2).
2010 *Mathematics Subject Classification*. Primary 47H09, 47J25, 90C25, 03F10

Among many other things, Bruck showed that in Hilbert spaces and for bounded closed and convex subsets C this iteration strongly converges for so-called acceptably paired sequences $(\lambda_n), (\theta_n)$ (cf. Definition 1.2). Moreover the limit is a fixed point of T provided that T is demicontinuous (continuous from the strong to the weak topology on H) in addition to being pseudocontractive:

Definition 1.2 ([1]). Two sequences (λ_n) and (θ_n) in $[0, 1]$ are **acceptably paired** if (θ_n) is nonincreasing, $\lim_{n \rightarrow \infty} \theta_n = 0$ and there exists a strictly increasing sequence $(f(n))_n$ of positive integers such that

1. $\liminf_{n \rightarrow \infty} \theta_{f(n)} \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j > 0,$
2. $\lim_{n \rightarrow \infty} (\theta_{f(n)} - \theta_{f(n+1)}) \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j = 0,$ and
3. $\lim_{n \rightarrow \infty} \sum_{j=f(n)}^{f(n+1)} \lambda_j^2 = 0.$

Theorem 1.3 (Corollary 4 of [1]). *Let C be a nonempty bounded closed convex subset of a Hilbert space H and $T : C \rightarrow C$ be a demicontinuous pseudocontraction. If (λ_n) and (θ_n) are acceptably paired such that $\lambda_n(1 + \theta_n) \leq 1$, then, for all $x_1, z \in C$, the sequence (x_n) defined by*

$$x_{n+1} = (1 - \lambda_n)x_n + \lambda_n T x_n + \lambda_n \theta_n(z - x_n)$$

remains in C and converges strongly to the fixed point of T which is closest to z .

Effective uniform rates on the strong convergence of (x_n) are generally ruled out. In fact, Neumann [8] showed that there are (computable) nonexpansive mappings f on the Hilbert cube (sequences $(x_n) \in \ell_2$ with $|x_n| \leq 1$ for all n) that have no computable fixed points, and so no sequence approximating any fixed point of f can have a computable rate of convergence. Following general proof-theoretic methods, it is necessary to pass first to a finitary version of Cauchyness, the so-called metastability in the sense of Tao, i.e. (here $[n; n + g(n)] := \{n, n + 1, n + 2, \dots, n + g(n)\}$)

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \in \mathbb{N} \forall i, j \in [n; n + g(n)] (\|x_i - x_j\| < \varepsilon).$$

Metastability is the so-called Herbrand normal form of (a suitable reformulation of) the Cauchy statement for the sequence (x_n) , and, as such, is equivalent to the original statement. It is finitary in the sense that it only talks about finite subsequences of (x_n) . A rate of metastability is then a bound $\Phi : \mathbb{N} \times \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}$ on the existential quantifier:

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Phi(\varepsilon, g) \forall i, j \in [n; n + g(n)] (\|x_i - x_j\| < \varepsilon).$$

Such bounds are guaranteed to exist under vastly general conditions on the complexity of the proof ([4]).

A quantitative, finitary version of all of Theorem 1.3, however, should not only finitise the Cauchyness of (x_n) , but also that the strong limit is indeed a fixed point. If T were norm-to-norm continuous, one way to do so would be to ensure that the sequence (x_n) is not only Cauchy along the interval $[n; n + g(n)]$, but also asymptotically regular:

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Phi(\varepsilon, g) \forall i, j \in [n; n + g(n)] (\|x_i - x_j\| < \varepsilon \wedge \|T x_i - x_i\| < \varepsilon).$$

By the logical equivalence of a statement to its Herbrand normal form, this implies both Cauchyness and asymptotic regularity. Cauchyness then implies that the strong limit exists, while norm-to-norm continuity and asymptotic regularity recover the fact that the limit is indeed a fixed point.

In the case at hand, however, the operator T is only demicontinuous. In fact, convergence to a fixed point is established via the *continuous* path (z_t) defined by $z_t = tT z_t + (1 - t)z$, which – in turn – converges strongly to the fixed point of T closest to z . This gives rise to the following finitary version of Theorem 1.3:

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Phi(\varepsilon, g) \forall i, j \in [n; n + g(n)] (\|x_i - x_j\| < \varepsilon \wedge \|x_i - y_i\| < \varepsilon \wedge \|y_i - T y_i\| < \varepsilon), (+)$$

where $y_i = z_{1/1+\theta_i}$. Our main theorem (Theorem 3.8) provides such a bound. If T is even norm-to-norm uniformly continuous with modulus ω , then one can obtain a bound Δ such that (see Theorem 3.11)

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Delta(g, \varepsilon) \forall i, j \in [n; n + g(n)] (\|x_i - x_j\| \leq \varepsilon \wedge \|x_i - Tx_i\| \leq \varepsilon).$$

This is a generalization of Theorem 2.8 of [6], which required T to be Lipschitz continuous. As guaranteed by general logical metatheorems [4], these bounds are highly uniform in the input data; it is independent of the space and the concrete choices for the operator $T : C \rightarrow C$, the set C or the parameter sequences (λ_n) and (θ_n) . Apart from the counterfunction g and the accuracy ε , the bounds only depend on an upper bound on the diameter $\text{diam}(C)$, moduli for the quantitative version of acceptably pariedness (cf. Definition 3.1) and, in the case of Theorem 3.11, the modulus of uniform continuity ω .

Moreover, the new, logically transformed proof of (+) is totally elementary in that all ideal principles have been eliminated; it can be formalized in constructive (“intuitionistic”) arithmetic enriched by an abstract normed space X (see Kohlenbach [4]) and axioms asserting that X is a Hilbert space. Moreover, one can recover Bruck’s original theorem using only the axiom of choice over *quantifier-free* sentences.

2 Analysis of Bruck’s Proof

We now examine from a proof-theoretic perspective the steps into which Bruck’s proof of Theorem 1.3 decomposes. First of all, we need to recall the generalization of pseudocontractiveness to set-valued operators. $T \subseteq H \times H$ is pseudocontractive, if, for all $(u, x), (v, y) \in T$,

$$\langle x - y, u - v \rangle \leq \|u - v\|^2.$$

Moreover, an operator $U \subset H \times H$ is monotone if and only if $I - U$ is pseudocontractive. It is *maximal* monotone if there does not exist a monotone $U' \subset H \times H$ such that $U \subsetneq U'$.

Bruck’s proof then follows the following line of argument:

- (i) The monotone operator $U : I - T$ is extended to a maximal monotone, set valued operator $U^* \subset H \times H$.
- (ii) There exists a unique y_θ for each $\theta > 0$ for which $0 \in \theta(y_\theta - z) + U^*(y_\theta)$.
- (iii) The strong $\lim_{\theta \rightarrow 0+} y_\theta$ exists and is the point x^* of $U^{*-1}(0)$ closest to z .
- (iv) The sequence (x_n) also converges to x^* .
- (v) The limit is a zero of U , and hence a fixed point of T .

The existence of a maximal monotone extension of a monotone operator $U : H \rightarrow H$ makes use of Zorn’s Lemma, which is equivalent to the Axiom of Choice. However, we are, for this paper, only interested in the single-valued case. As shown in [6], it is possible to avoid the detour via maximal monotone extensions. A similar result has also been shown by Lan and Wu in [7]. The existence of the path $(y_t)_{t \in (0,1]}$ is also guaranteed in this case since the mapping $U_t : C \rightarrow C, y \mapsto t(y - z) + U(x)$ is t -strongly monotone for each $t > 0$, and thus has a unique fixed point (see [6]). The mere existence of the sequence $(y_{\theta_n})_n$ makes no proof-theoretic contribution since their defining property is a purely universal statement, i.e. one with only $\forall$ -quantifiers.

The convergence of (y_{θ_n}) to the fixed point of T closest to z is then carried out analogously to the multi-valued case in Bruck’s proof [1]. A quantitative analysis of this step has been performed and a rate of metastability has already been extracted in [6].

The convergence $\|x_n - x^*\| \rightarrow 0$ is established via convergence of the subsequence $\|x_{f(n)} - x^*\| \rightarrow 0$, which is shown using the existence of the limit superior as a translation invariant functional $\limsup : \ell_\infty \rightarrow \mathbb{R}$ as follows: If $f : \mathbb{N} \rightarrow \mathbb{N}$ denotes the subsequence from Definition 1.2, then there exists a constant $\gamma \in (0, 1)$ such that

$$\begin{aligned} \gamma \cdot \limsup \|x_{f(k)} - x^*\|^2 &= \gamma \cdot \limsup \|x_{f(k)} - y_{\theta_{f(k)}}\|^2 \\ &\geq \limsup \|x_{f(k+1)} - y_{\theta_{f(k)}}\|^2 \\ &= \limsup \|x_{f(k+1)} - x^*\|^2 \\ &= \limsup \|x_{f(k)} - x^*\|^2. \end{aligned} \tag{1}$$

where inequality (1) is shown in Bruck's proof. Therefore, $\limsup \|x_{f(k)} - x^*\| = 0$, so the subsequence $(x_{f(k)})$ converges to x^* . Basic arithmetic then implies the convergence of the original sequence.

3 Main Results

To obtain a quantitative version of Theorem 1.3, we need a quantitative version of what it means for two sequences to be acceptably paired.

Definition 3.1. Two sequences (λ_n) and (θ_n) in $[0, 1]$ are called acceptably paired with moduli $\varphi_1, \varphi_2, \varphi_3 : \mathbb{R} \rightarrow \mathbb{N}$, $f : \mathbb{N} \rightarrow \mathbb{N}$, $n_0 \in \mathbb{N}$ and $\delta > 0$ if (θ_n) is nonincreasing and the following conditions are satisfied:

1. $\forall \varepsilon > 0 \forall n \geq \varphi_1(\varepsilon) (\theta_n \leq \varepsilon)$,
2. $\forall n (f(n+1) \geq f(n) + 1)$,
3. $\forall n \geq n_0 \left(\theta_{f(n)} \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j \geq \delta \right)$,
4. $\forall \varepsilon > 0 \forall n \geq \varphi_2(\varepsilon) \left((\theta_{f(n)} - \theta_{f(n+1)}) \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j \leq \varepsilon \right)$, and
5. $\forall \varepsilon > 0 \forall n \geq \varphi_3(\varepsilon) \left(\sum_{j=f(n)}^{f(n+1)} \lambda_j^2 \leq \varepsilon \right)$.

The moduli φ_i are rates of convergence of their respective sequences to 0. The numbers n_0 and δ are quantitative witnesses for the condition that the sequence $\theta_{f(n)} \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j$ stays strictly away from 0, i.e. its $\liminf$ is greater than 0. It is also noteworthy that the function $k \mapsto k^* := \max\{n \in \mathbb{N} : f(n) \leq k\}$ is well-defined for all $k \geq f(0)$. Moreover, $(f(k))^* = k$ for all nonnegative integers k .

Remark 3.2 ([1]). Examples of acceptably paired sequences are:

1. $\lambda_n = 1/n$, $\theta_n = 1/\log \log n$ and $f(n) = n^n$.
2. For $0 < p < 1$ and $0 < q < \min\{p, 1-p\}$, $\lambda_n = n^{-p}$ and $\theta_n = n^{-q}$ are acceptably paired with $f(n) = \lceil n^{d/(1-p)} \rceil$ for suitable $d > 1$ (see Section 4.1 for details).

The corresponding moduli will be given in Section 4.

Lemma 3.3. Suppose that X is a normed space and $(a_n) \subseteq X$ is metastable with rate $\Psi : (0, \infty) \times \mathbb{N} \rightarrow \mathbb{N}$. Then, for any nondecreasing $f : \mathbb{N} \rightarrow \mathbb{N}$ with $f(n) \geq n$, the sequence $(a_{f(n)})$ is metastable with rate $\tilde{\Psi}_f$ defined by $\tilde{\Psi}_f(\varepsilon, g) := \Psi(\varepsilon, g_f)$, where $g_f : \mathbb{N} \rightarrow \mathbb{N}$ is defined by $g_f(n) := f(n + g(n)) - n$.

Proof. Since (a_n) is metastable with modulus Ψ ,

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Psi(\varepsilon, g_f) \forall i, j \in [n; f(n + g(n))] (\|a_i - a_j\| \leq \varepsilon).$$

Since $f(n) \geq n$, we conclude

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Psi(\varepsilon, g_f) \forall i, j \in [f(n); f(n + g(n))] (\|a_i - a_j\| \leq \varepsilon).$$

The monotonicity of f then implies

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Psi(\varepsilon, g_f) \forall i, j \in [n; n + g(n)] (\|a_{f(i)} - a_{f(j)}\| \leq \varepsilon),$$

so $\tilde{\Psi}_f$ is a rate of metastability for $(a_{f(n)})$. □

Lemma 3.4. Suppose that $f : \mathbb{N} \rightarrow \mathbb{N}$ is strictly increasing and for each k , we have a statement $A(k)$. Define a function $(\cdot)^* : \{n \in \mathbb{N} : n \geq f(0)\} \rightarrow \mathbb{N}$ by $k \mapsto \max\{n \in \mathbb{N} : f(n) \leq k\}$. Then for all $g : \mathbb{N} \rightarrow \mathbb{N}$

$$A(k) \text{ for all } k \in [n; n + \tilde{g}(n)] \Rightarrow A(k^*) \text{ for all } k \in [m; m + g(m)],$$

where $\tilde{g}(n) := (f(n) + g(f(n)))^* - n$ and $m := f(n)$.

Proof. Assume the statement $A(k)$ holds for all $k \in [n; n + \tilde{g}(n)]$. Observe that $n + \tilde{g}(n) = (m + g(m))^*$ and $n = (f(n))^* = m^*$, so the statement $A(k)$ holds for all $k \in [m^*; (m + g(m))^*]$. Therefore,

$$A(m^*) \wedge A(m^* + 1) \wedge \dots \wedge A((m + g(m))^*)$$

In particular, $(\cdot)^*$ is nondecreasing (since f is nondecreasing) and so

$$A(m^*) \wedge A((m + 1)^*) \wedge \dots \wedge A((m + g(m))^*).$$

Therefore, statement $A(k^*)$ holds for all $k \in [m; m + g(m)]$. $\square$

We now give our main results, which were obtained by logical analysis of Bruck's proof [1] using the proof-theoretic methods treated extensively in [4].

Theorem 3.5. *Let C be a nonempty bounded closed convex subset of a Hilbert space H with $\text{diam}(C) \leq M \in \mathbb{N}$, $T : C \rightarrow C$ be a demicontinuous, single-valued pseudocontraction and $x_1, z \in C$. Suppose the sequences (λ_n) and (θ_n) are acceptably paired with moduli as in Definition 3.1 satisfying $\lambda_n(1 + \theta_n) \leq 1$, and the sequence (y_i) defined by*

$$y_i = \frac{1}{1 + \theta_i} T y_i + \frac{\theta_i}{1 + \theta_i} z$$

is metastable with rate $\Psi : (0, \infty) \times \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}$. Define the sequence (x_n) by

$$x_{n+1} = (1 - \lambda_n) x_n + \lambda_n T x_n + \lambda_n \theta_n (z - x_n),$$

and a function Φ by $\Phi(\varepsilon, g, \varphi_1, \varphi_2, \varphi_3, \delta, n_0, M, f) := f(\tilde{\Psi}(\tilde{\varepsilon}, g_d) + n_1 + d + 1)$, where $\varphi_1, \varphi_2, \varphi_3 : \mathbb{R} \rightarrow \mathbb{N}$, $f : \mathbb{N} \rightarrow \mathbb{N}$, $n_0 \in \mathbb{N}$ and $\delta > 0$ are the moduli of Definition 3.1 and

$$\begin{aligned} \tilde{\Psi}_f(\varepsilon, g) &:= \Psi(\varepsilon, g_f), & g_f(n) &:= f(n + g(n)) - n \\ g_d(n) &:= d + n_1 + 1 + \tilde{g}(n + n_1 + d + 1) & \tilde{g}(n) &:= (f(n) + g(f(n)))^* - n \\ d &:= \max\{f(k_0), \lceil \log_c(\varepsilon/8M) \rceil\}, & k^* &:= \max\{n \in \mathbb{N} : f(n) \leq k\} \\ n_1 &:= \max\{n_0, \varphi_1(\delta/2), \varphi_2(\tilde{\varepsilon}/4M^2), \varphi_3(\tilde{\varepsilon}^2/8M^2)\}, & k_0 &:= \max\{\varphi_2(\varepsilon^2/6M^2), \varphi_3(\varepsilon^2/12M^2)\}, \\ c &:= \exp(-\delta/2)m, & \tilde{\varepsilon} &:= \frac{1-c}{16} \cdot \varepsilon. \end{aligned}$$

To simplify notation, we will omit the dependence of Φ on the moduli for the parameters (λ_n) and (θ_n) and instead write $\Phi(\varepsilon, g) := \Phi(\varepsilon, g, \varphi_1, \varphi_2, \varphi_3, \delta, n_0, M, f)$. Then,

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Phi(g, \varepsilon) \forall i, j \in [n; n + g(n)] (\|x_i - y_{f(i^*)}\| \leq \varepsilon \wedge \|y_{f(i^*)} - y_{f(j^*)}\| \leq \varepsilon)$$

Remark 3.6. Observe that the bound given in Theorem 3.5 is independent of the operator T and the space H . Moreover, it is also highly uniform with respect to the domain C (dependence only via an upper bound on the diameter $\text{diam} C$) and the choice of the parameter sequences (λ_n) and (θ_n) (dependence only via the moduli $\varphi_1, \varphi_2, \varphi_3, \delta, n_0$ and f).

Proof. Since T is pseudocontractive, $U := I - T$ is monotone. Moreover,

$$\begin{aligned} y_i &= \frac{1}{1 + \theta_i} T y_i + \frac{\theta_i}{1 + \theta_i} z \\ &= \frac{1}{1 + \theta_i} (I - U)(y_i) + \frac{\theta_i}{1 + \theta_i} z \\ &= \frac{1}{1 + \theta_i} y_i - \frac{1}{1 + \theta_i} U y_i + \frac{\theta_i}{1 + \theta_i} z, \end{aligned}$$

which is equivalent to

$$0 = \left(1 - \frac{1}{1 + \theta_i}\right) y_i + \frac{1}{1 + \theta_i} U y_i - \frac{\theta_i}{1 + \theta_i} z = \frac{\theta_i}{1 + \theta_i} (y_i - z) + \frac{1}{1 + \theta_i} U y_i,$$

so $\theta_i(y_i - z) + Uy_i = 0$. Moreover, the Bruck iteration rewritten in terms of U reads

$$\begin{aligned} x_{n+1} &= (1 - \lambda_n)x_n + \lambda_n T x_n + \lambda_n \theta_n(z - x_n) \\ &= x_n - \lambda_n(x_n - T x_n + \theta_n(x_n - z)) \\ &= x_n - \lambda_n(Ux_n + \theta_n(x_n - z)). \end{aligned}$$

Therefore, for $n > i \geq 2$,

$$x_n - y_i = x_{n-1} - y_i - \lambda_{n-1}(Ux_{n-1} + \theta_{n-1}(x_{n-1} - z)),$$

so

$$\begin{aligned} \|x_n - y_i\|^2 &= \left\langle x_{n-1} - y_i - \lambda_{n-1}(Ux_{n-1} + \theta_{n-1}(x_{n-1} - z)), \right. \\ &\quad \left. x_{n-1} - y_i - \lambda_{n-1}(Ux_{n-1} + \theta_{n-1}(x_{n-1} - z)) \right\rangle \\ &= \|x_{n-1} - y_i\|^2 - 2\lambda_{n-1}\langle x_{n-1} - y_i, Ux_{n-1} + \theta_{n-1}(x_{n-1} - z) \rangle \\ &\quad + \lambda_{n-1}^2 \|Ux_{n-1} + \theta_{n-1}(x_{n-1} - z)\|^2 \\ &= \|x_{n-1} - y_i\|^2 + \lambda_{n-1}^2 \|Ux_{n-1} + \theta_{n-1}(x_{n-1} - z)\|^2 \\ &\quad - 2\lambda_{n-1}\theta_{n-1}\langle x_{n-1} - y_i, x_{n-1} - z \rangle \\ &\quad - 2\lambda_{n-1}\langle x_{n-1} - y_i, Ux_{n-1} \rangle \\ &= \|x_{n-1} - y_i\|^2 + \lambda_{n-1}^2 \|Ux_{n-1} + \theta_{n-1}(x_{n-1} - z)\|^2 \\ &\quad + 2\lambda_{n-1}(\theta_i - \theta_{n-1})\langle x_{n-1} - y_i, x_{n-1} - z \rangle \\ &\quad - 2\lambda_{n-1}\langle x_{n-1} - y_i, Ux_{n-1} + \theta_i(x_{n-1} - z) \rangle. \end{aligned} \tag{2}$$

Since U is monotone and $\theta_i(y_i - z) + Uy_i = 0$,

$$\begin{aligned} \langle Ux_{n-1} + \theta_i(x_{n-1} - z), x_{n-1} - y_i \rangle &= \langle Ux_{n-1} + \theta_i(y_i - z), x_{n-1} - y_i \rangle + \theta_i \|x_{n-1} - y_i\|^2 \\ &= \langle Ux_{n-1} - Uy_i, x_{n-1} - y_i \rangle + \theta_i \|x_{n-1} - y_i\|^2 \\ &\geq \theta_i \|x_{n-1} - y_i\|^2. \end{aligned}$$

Equation (2) then implies

$$\begin{aligned} \|x_n - y_i\|^2 &\leq (1 - 2\lambda_{n-1}\theta_i) \|x_{n-1} - y_i\|^2 + \lambda_{n-1}^2 \|Ux_{n-1} + \theta_{n-1}(x_{n-1} - z)\|^2 \\ &\quad + 2\lambda_{n-1}(\theta_i - \theta_{n-1})\langle x_{n-1} - z, x_{n-1} - y_i \rangle. \end{aligned}$$

Observe that $\|Ux_{n-1} + \theta_{n-1}(x_{n-1} - z)\| = \|x_{n-1} - Tx_{n-1} + \theta_{n-1}(x_{n-1} - z)\|$. Since $\text{diam}(C) \leq M$, we conclude

$$\|x_n - y_i\|^2 \leq \exp(-2\lambda_{n-1}\theta_i) \|x_{n-1} - y_i\|^2 + 2M^2\lambda_{n-1}(\theta_i - \theta_{n-1}) + 4M^2\lambda_{n-1}^2. \tag{3}$$

We show by induction on $n \geq i$ that

$$\|x_n - y_i\|^2 \leq \exp\left(-2\theta_k \sum_{j=i}^{n-1} \lambda_j\right) \|x_i - y_i\|^2 + 2M^2 \sum_{j=i}^{n-1} (\theta_i - \theta_j) \lambda_j + 4M^2 \sum_{j=i}^{n-1} \lambda_j^2. \tag{4}$$

Proof of (4): For $n = i$ the inequality holds with equality. Suppose that the inequality holds true for

some $n \geq i$. Then (3) implies

$$\begin{aligned}
\|x_{n+1} - y_i\|^2 &\leq \exp(-2\lambda_n \theta_i) \|x_n - y_i\|^2 + 2M^2 \lambda_n (\theta_i - \theta_n) + 4M^2 \lambda_n^2 \\
&\leq \exp(-2\lambda_n \theta_i) \cdot \left\{ \exp\left(-2\theta_i \sum_{j=i}^{n-1} \lambda_j\right) \|x_i - y_i\|^2 \right. \\
&\quad \left. + 2M^2 \sum_{j=i}^{n-1} (\theta_i - \theta_j) \lambda_j + 4M^2 \sum_{j=i}^{n-1} \lambda_j^2 \right\} \\
&\quad + 2M^2 \lambda_n (\theta_i - \theta_n) + 4M^2 \lambda_n^2 \\
&= \exp\left(-2\theta_i \sum_{j=i}^n \lambda_j\right) \|x_i - y_i\|^2 + 2M^2 \sum_{j=i}^n (\theta_i - \theta_j) \lambda_j + 4M^2 \sum_{j=i}^n \lambda_j^2,
\end{aligned}$$

which is what we needed to show.

Since $\theta_i - \theta_j \leq \theta_i - \theta_n$ for $i \leq j \leq n$, (4) implies

$$\|x_n - y_i\|^2 \leq \exp\left(-2\theta_i \sum_{j=i}^{n-1} \lambda_j\right) \|x_i - y_i\|^2 + 2M^2 (\theta_i - \theta_n) \sum_{j=i}^{n-1} \lambda_j + 4M^2 \sum_{j=i}^{n-1} \lambda_j^2, \text{ for all } n \geq i. \quad (5)$$

Now let $f(n)$ be the subsequence of Definition 1.2. We now prove that $(x_{f(n)})$ is Cauchy. Taking $i = f(k)$ and $n = f(k+1)$ in (5), we get

$$\begin{aligned}
\|x_{f(k+1)} - y_{f(k)}\|^2 &\leq \exp\left(-2\theta_{f(k)} \sum_{j=f(k)}^{f(k+1)} \lambda_j\right) \cdot \exp(2\theta_{f(k)} \lambda_{f(k+1)}) \cdot \|x_{f(k)} - y_{f(k)}\|^2 \\
&\quad + 2M^2 (\theta_{f(k)} - \theta_{f(k+1)}) \cdot \sum_{j=f(k)}^{f(k+1)} \lambda_j + 4M^2 \sum_{j=f(k)}^{f(k+1)} \lambda_j^2. \quad (6)
\end{aligned}$$

Now observe that

$$\exp\left(-2\theta_{f(k)} \sum_{j=f(k)}^{f(k+1)} \lambda_j\right) \leq \exp(-2\delta) < 1, \text{ for all } k \geq n_0.$$

Moreover, (θ_n) is a null sequence with modulus φ_1 . Thus, $\exp(2\theta_{f(k)} \lambda_{f(k+1)}) \leq \exp(2\theta_k) \leq \exp(\delta)$ for all $k \geq \varphi_1(\delta/2)$. Furthermore, for all $k \geq \max\{\varphi_2(\tilde{\varepsilon}^2/4M^2), \varphi_3(\tilde{\varepsilon}^2/8M^2)\}$, the remainder term in (6) is less than $\tilde{\varepsilon}^2$. In total,

$$\|x_{f(k+1)} - y_{f(k)}\|^2 \leq \exp(-\delta) \cdot \|x_{f(k)} - y_{f(k)}\|^2 + \tilde{\varepsilon}^2, \text{ for all } k \geq n_1$$

since $n_1 = \max\{n_0, \varphi_1(\delta/2), \varphi_2(\tilde{\varepsilon}^2/4M^2), \varphi_3(\tilde{\varepsilon}^2/8M^2)\}$. Because $c = \exp(-\delta/2)$, we then get

$$\begin{aligned}
\|x_{f(k+1)} - y_{f(k)}\| &\leq c \cdot \|x_{f(k)} - y_{f(k)}\| + \tilde{\varepsilon} \\
&\leq c \cdot \|x_{f(k)} - y_{f(k-1)}\| + c \cdot \|y_{f(k-1)} - y_{f(k)}\| + \tilde{\varepsilon}. \quad (7)
\end{aligned}$$

Now observe that since (y_n) is metastable with rate Ψ , the subsequence $(y_{f(n)})$ is metastable with rate $\tilde{\Psi}$ by Lemma 3.3. Thus, there exists an integer $n \leq \tilde{\Psi}(\tilde{\varepsilon}, g_d)$ such that $\|y_{f(k)} - y_{f(j)}\| \leq \tilde{\varepsilon}$ for all $k, j \in [n; n + g_d(n)]$. Taking $n_2 := n + n_1$, we have on the one hand $n_2 \geq n_1$, and $\|y_{f(k)} - y_{f(j)}\| \leq \tilde{\varepsilon}$ for all $k, j \in [n_2; n_2 + d + 1 + \tilde{g}(n_2 + d + 1)]$ on the other. Setting $j = k - 1$, we conclude

$$\|y_{f(k)} - y_{f(k-1)}\| \leq \tilde{\varepsilon}, \text{ for all } k \in [n_2 + 1; n_2 + d + 1 + \tilde{g}(n_2 + d + 1)]. \quad (8)$$

Suppose now that $k \in [n_2 + d; n_2 + d + \tilde{g}(n_2 + d + 1)]$. Then (8) and (7) yield

$$\begin{aligned}
\|x_{f(k+1)} - y_{f(k)}\| &\leq c \cdot \|x_{f(k)} - y_{f(k-1)}\| + 2\tilde{\varepsilon} \\
&\leq c \cdot (c \cdot \|x_{f(k-1)} - y_{f(k-2)}\| + 2\tilde{\varepsilon}) + 2\tilde{\varepsilon} \\
&= c^2 \cdot \|x_{f(k-1)} - y_{f(k-2)}\| + 2\tilde{\varepsilon} \cdot c + 2\tilde{\varepsilon} \\
&\leq \dots \\
&= c^{d-1} \cdot \|x_{f(k-d+2)} - y_{f(k-d+1)}\| + 2\tilde{\varepsilon} \sum_{k=0}^{d-2} c^k \\
&\leq c^d \cdot \|x_{f(k-d+1)} - y_{f(k-d)}\| + c^{d-1} \cdot \|y_{f(k-d+1)} - y_{f(k-d)}\| + 2\tilde{\varepsilon} \sum_{k=0}^{d-2} c^k \\
&\leq c^d \cdot M + 2\tilde{\varepsilon} \sum_{k=0}^{d-1} c^k \\
&\leq c^d \cdot M + 2\tilde{\varepsilon} \sum_{k=0}^{\infty} c^k \\
&= c^d \cdot M + \frac{2\tilde{\varepsilon}}{1-c}.
\end{aligned}$$

Since $d \geq \log_c(\varepsilon/8M)$ and $\tilde{\varepsilon} = \frac{1-c}{16} \cdot \varepsilon$, we have

$$\|x_{f(k+1)} - y_{f(k)}\| \leq \frac{\varepsilon}{4}, \text{ for all } k \in [n_2 + d; n_2 + d + \tilde{g}(n_2 + d + 1)].$$

Therefore, setting $n_3 := n_2 + d + 1$ and using (8),

$$\begin{aligned}
\|x_{f(k)} - y_{f(k)}\| &\leq \|x_{f(k)} - y_{f(k-1)}\| + \|y_{f(k)} - y_{f(k-1)}\| \leq \frac{\varepsilon}{4} + \tilde{\varepsilon} \\
&< \frac{\varepsilon}{3}, \text{ for all } k \in [n_3; n_3 + \tilde{g}(n_3)].
\end{aligned}$$

By Lemma 3.4

$$\|x_{f(k^*)} - y_{f(k^*)}\| \leq \varepsilon/3, \text{ for all } k \in [f(n_3); f(n_3) + g(f(n_3))] \quad (9)$$

Now, for $k \geq f(0)$, observe that k^* denotes the unique integer such that $f(k^*) \leq k < f(k^* + 1)$. Take $n = k$, $i = f(k^*)$ in (5); since the exponential factor is less than or equal to 1,

$$\begin{aligned}
\|x_k - y_{f(k^*)}\|^2 &\leq \|x_{f(k^*)} - y_{f(k^*)}\|^2 + 2M^2(\theta_{f(k^*)} - \theta_k) \sum_{j=f(k^*)}^{k-1} \lambda_j + 4M^2 \sum_{j=f(k^*)}^{k-1} \lambda_j^2 \\
&\leq \|x_{f(k^*)} - y_{f(k^*)}\|^2 + 2M^2(\theta_{f(k^*)} - \theta_{f(k^*+1)}) \sum_{j=f(k^*)}^{f(k^*+1)} \lambda_j + 4M^2 \sum_{j=f(k^*)}^{f(k^*+1)} \lambda_j^2.
\end{aligned}$$

Observe that the latter two terms become less than $\varepsilon^2/3$ whenever $k^* \geq k_0$ since, by definition, $k_0 = \max\{\varphi_2(\varepsilon^2/6M^2), \varphi_3(\varepsilon^2/12M^2)\}$. But this is always the case whenever $k \geq f(k_0)$ since then $k^* \geq (f(k_0))^* = k_0$ by the monotonicity of $(\cdot)^*$. Therefore,

$$\|x_k - y_{f(k^*)}\|^2 \leq \|x_{f(k^*)} - y_{f(k^*)}\|^2 + \frac{2\varepsilon^2}{3}, \text{ for all } k \geq f(k_0). \quad (10)$$

Since $f(n_3) = f(n_2 + d + 1) \geq f(d) \geq d \geq f(k_0)$, equations (9), (10) together imply

$$\|x_k - y_{f(k^*)}\| \leq \varepsilon, \text{ for all } k \in [f(n_3); f(n_3) + g(f(n_3))]. \quad (11)$$

Now recall that $\|y_{f(i)} - y_{f(j)}\| \leq \tilde{\varepsilon}$ for all $i, j \in [n_2; n_2 + d + 1 + \tilde{g}(n_2 + d + 1)]$. Again by Lemma 3.4, this implies

$$\|y_{f(i^*)} - y_{f(j^*)}\| \leq \tilde{\varepsilon} \leq \varepsilon, \text{ for all } i, j \in [f(n_3), f(n_3) + g(f(n_3))]. \quad (12)$$

Therefore, $f(n_3) = f(n_2 + d + 1) \leq f(\tilde{\Psi}(g_d, \tilde{\varepsilon}) + n_1 + d + 1)$ satisfies the claim. $\square$

Theorem 3.7. *In the situation of Theorem 3.5, (x_n) is metastable with rate $\Phi'(\varepsilon, g) := \Phi(\varepsilon/3, g)$.*

Proof. Since $\tilde{\varepsilon} < \varepsilon$ and $\|x_i - x_j\| \leq \|x_i - y_{f(i^*)}\| + \|x_j - y_{f(j^*)}\| + \|y_{f(i^*)} - y_{f(j^*)}\|$, equations (11) and (12) imply

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Phi(\varepsilon/3, g) \forall i, j \in [n; n + g(n)] (\|x_i - x_j\| < \varepsilon),$$

which is what we needed to show. $\square$

Theorem 3.8. *In the situation of Theorem 3.5,*

$$\begin{aligned} \forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Phi''(\varepsilon, g) \forall i, j \in [n; n + g(n)] \\ (\|x_i - x_j\| < \varepsilon \wedge \|x_i - y_i\| < \varepsilon \wedge \|y_i - Ty_i\| < \varepsilon), \end{aligned}$$

where $\Phi''(\varepsilon, g) := \hat{\Phi}(\varepsilon/3,)$, and $\hat{\Phi}$ is defined like Φ in Theorem 3.5, but with $\hat{g}_d$, defined by

$$\hat{g}_d(n) := d + \hat{n}_1 + 1 + \hat{g}(n + d + \hat{n}_1 + 1).$$

instead of g_d , where $\hat{g}(n) := (f(n) + g(f(n)))^* - n + 1$ and $\hat{n}_1 := \max\{n_1, \varphi_1(\varepsilon/M)\}$.

Moreover, we can take $\Psi(\varepsilon, g) := \tilde{g}^{(\lceil 16d^2/\varepsilon^2 \rceil)}(1)$, where $\tilde{g}(n) = n + 1 + g(n + 1)$.

Proof. By altering the definition of g_d , the point $f(n_3)$ that satisfies the conclusion of Theorem 3.7 also satisfies $f(n_3) \geq n_3 = n_2 + d + 1 = n_0 + \hat{n}_1 + d + 1 \geq \varphi_1(\varepsilon, g)$. Therefore, (x_n) is metastable, and

$$\|y_i - Ty_i\| = \frac{\theta_i}{1 + \theta_i} \|Ty_i - z\| \leq \theta_i \cdot M \leq \varepsilon, \text{ for all } i \geq n.$$

It remains to verify that $\|x_i - y_i\| \leq \varepsilon$ on $[f(n_3); f(n_3) + g(f(n_3))]$. To this end, observe that

$$\|y_i - y_j\| \leq \varepsilon/3, \text{ for all } i, j \in [f(n_3); f(n_3) + \hat{g}(n_3)]. \quad (13)$$

Since $f(k^* + 1) > k \geq f(k^*)$ for all $k \geq f(0)$, we conclude $f(n_3 + \hat{g}(n_3)) = f((f(n_3) + g(f(n_3)))^* + 1) \geq f(n_3) + g(f(n_3))$. Moreover, $f((f(n_3))^*) = f(n_3)$. Therefore, (13) implies

$$\|y_k - y_{f(k^*)}\| \leq \varepsilon/3, \text{ for all } k \in [f(n_3), f(n_3) + g(f(n_3))],$$

which, using (11), implies for all $k \in [f(n_3), f(n_3) + g(f(n_3))]$

$$\|x_k - y_k\| \leq \|x_k - y_{f(k^*)}\| + \|y_{f(k^*)} - y_k\| \leq \varepsilon/3 + \varepsilon/3 < \varepsilon.$$

That we may choose $\Psi(\varepsilon, g) := \tilde{g}^{(\lceil 16d^2/\varepsilon^2 \rceil)}(1)$ follows from Theorem 2.8 and Corollary 2.9 of [6]. $\square$

Remark 3.9. Observe that Theorems 3.5, 3.7 and 3.8 require only the demicontinuity of T . Therefore, model-theoretic approaches (cf. [3]) are not applicable, as these always require norm-continuity.

Remark 3.10. Suppose Ψ does not depend on g for a concrete choice of the input. Then metastability for (y_n) would read

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Psi(\varepsilon) \forall i, j \in [n; n + g(n)] (\|x_i - x_j\| < \varepsilon).$$

This is logically equivalent to

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Psi(\varepsilon) \forall i, j \geq n (\|x_i - x_j\| < \varepsilon),$$

i.e. a rate of convergence. In this case, we would get in Theorem 3.5 a rate of convergence $\Phi(\varepsilon) := f(\Psi(\tilde{\varepsilon}) + n_1 + d + 1)$, where, as before

$$\begin{aligned} n_1 &:= \max\{n_0, \varphi_1(\delta/2), \varphi_2(\tilde{\varepsilon}/4M^2), \varphi_3(\tilde{\varepsilon}^2/8M^2)\}, & k_0 &:= \max\{\varphi_2(\varepsilon^2/6M^2), \varphi_3(\varepsilon^2/12M^2)\}, \\ d &:= \max\{f(k_0), \lceil \log_c(\varepsilon/8M) \rceil\}, & \tilde{\varepsilon} &= \frac{1-c}{16} \cdot \varepsilon, \\ c &:= \exp(-\delta/2). \end{aligned}$$

Theorem 3.11. *Suppose that in the situation of Theorem 3.5, T is additionally uniformly continuous on C with modulus ω . For $g : \mathbb{N} \rightarrow \mathbb{N}$ define $g_b(n) := b + g(n + b)$, where $b := f((\varphi_1(\varepsilon/3M))^* + 1)$. Then*

$$\forall \varepsilon > 0 \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists n \leq \Delta(g, \varepsilon) \forall i, j \in [n; n + g(n)] (\|x_i - x_j\| \leq \varepsilon \wedge \|x_i - Tx_i\| \leq \varepsilon),$$

where $\Delta(g, \varepsilon) := \Phi(g_b, \min\{\varepsilon/3, \omega(\varepsilon/3M)\}) + b$.

Proof. By Theorem 3.5, there exists a $k \leq \Phi(g_b, \min\{\varepsilon/3, \omega(\varepsilon/3M)\})$ such that for $n := k + b$,

$$\|x_i - y_{f(i^*)}\| \leq \min\{\varepsilon/3, \omega(\varepsilon/3M)\}, \text{ for all } i, j \in [n; n + g(n)].$$

Now observe that $b^* = (\varphi_1(\varepsilon/3M))^* + 1$ since $(f(k))^* = k$ for all nonnegative integers k . Therefore, $f(n^*) \geq f(b^*) = f((\varphi_1(\varepsilon/3M))^* + 1) > \varphi_1(\varepsilon/3M)$. Thus, $\theta_{f(i^*)} \leq \varepsilon/3M$ for all $i \geq n$. Consequently

$$\|y_{f(i^*)} - Ty_{f(i^*)}\| = \frac{\theta_{f(i^*)}}{1 + \theta_{f(i^*)}} \|Ty_{f(i^*)} - z\| \leq \theta_{f(i^*)} \cdot M \leq \frac{\varepsilon}{3}, \text{ for all } i \geq n.$$

Therefore,

$$\|x_i - Tx_i\| \leq \|x_i - y_{f(i^*)}\| + \|y_{f(i^*)} - Ty_{f(i^*)}\| + \|Tx_i - Ty_{f(i^*)}\| \leq \varepsilon, \text{ for all } i \in [n; n + g(n)].$$

That $\|x_i - x_j\| \leq \varepsilon$ on $[n; n + g(n)]$ follows as in Corollary 3.7. $\square$

4 Application to Concrete Instances

In this section, we compute explicitly the moduli $\varphi_1, \varphi_2, \varphi_3, n_0$ and δ for the two examples of parameter sequences of Remark 3.2. We then compare the bound to the one obtained in [5] for Halpern iterations of nonexpansive mappings.

4.1 Example 1

Suppose p and q are real numbers in $(0, 1)$ such that $0 < q < \min\{p, 1 - p\}$, and take $\lambda_n := n^{-p}$ and $\theta_n := n^{-q}$. Set $r = (p + q)/2$. There are two cases to consider, namely $p \geq 1/2$ and $p < 1/2$.

If $p < 1/2$, then $1/2 > p > r > q$, so we conclude $1/2 < 1 - p < 1 - r < 1 - q$, whence $0 < 1 - 2p < 1 - r - p < 1 - p - q$. Then,

$$\frac{1 - 2p}{1 - p} < \frac{1 - r - p}{1 - p} < \frac{1 - p - q}{1 - p} = 1 - \frac{q}{1 - p},$$

and so

$$\frac{1 - p}{1 - 2p} > \frac{1 - p}{1 - r - p} > \left(1 - \frac{q}{1 - p}\right)^{-1}.$$

Thus, if we choose $d := \min\left\{\frac{1 - p}{1 - r - p}, \frac{3}{2} \left(1 - \frac{q}{1 - p}\right)^{-1}\right\}$, then

$$\left(1 - \frac{q}{1 - p}\right)^{-1} < d < \min\left\{\frac{1 - p}{1 - 2p}, 2 \left(1 - \frac{q}{1 - p}\right)^{-1}\right\} \quad (\text{for } p < 1/2),$$

which is Bruck's condition. For $p \geq \frac{1}{2}$, we see as before that

$$\left(1 - \frac{q}{1 - p}\right)^{-1} < d < 2 \left(1 - \frac{q}{1 - p}\right)^{-1} \quad (\text{for } p \geq 1/2).$$

An important consequence of our choice of d is that $d > 1$, which we will use throughout this section.

Now, one can take $f(n) := \lceil n^{d/(1-p)} \rceil$. To calculate the other moduli, we need the following Lemma, which is a direct consequence of Taylor's Theorem using the Lagrange remainder term.

Lemma 4.1. Suppose $x, r \in \mathbb{R}$ with $x > 0$ and $r \neq 1$. Then,

- (i) there exists a real number $\xi \in (x, x+1)$ such that $(x+1)^r = x^r + r\xi^{r-1}$, and
- (ii) there exists a real number $\nu \in (x-1, x)$ such that $(x-1)^r = x^r - r\nu^{r-1}$.

We now proceed to calculate the moduli. Observe that

$$\begin{aligned}
\theta_{f(n)} \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j &= \left[n^{d/(1-p)} \right]^{-q} \cdot \sum_{j=f(n)}^{f(n+1)} j^{-p} \\
&> (n^{d/(1-p)} + 1)^{-q} \cdot \int_{f(n)}^{f(n+1)} j^{-p} dj \\
&= \frac{(n^{d/(1-p)} + 1)^{-q}}{1-p} \left[(f(n+1))^{1-p} - (f(n))^{1-p} \right] \\
&= \frac{(n^{d/(1-p)} + 1)^{-q}}{1-p} \left(\left[(n+1)^{d/(1-p)} \right]^{1-p} - \left[n^{d/(1-p)} \right]^{1-p} \right) \\
&\geq \frac{(n^{d/(1-p)} + 1)^{-q}}{1-p} \left[\left((n+1)^{d/(1-p)} \right)^{1-p} - \left(n^{d/(1-p)} + 1 \right)^{1-p} \right] \\
&= \frac{(n^{d/(1-p)} + 1)^{-q}}{1-p} \left[(n+1)^d - \left(n^{d/(1-p)} + 1 \right)^{1-p} \right]
\end{aligned} \tag{14}$$

By virtue of 4.1, there exists a $\xi \in (n^{d/(1-p)}, n^{d/(1-p)} + 1)$ such that

$$\begin{aligned}
\left(n^{d/(1-p)} + 1 \right)^{1-p} &= n^d + (1-p)\xi^{-p} \\
&\leq n^d + (1-p)n^{-\frac{dp}{1-p}}.
\end{aligned}$$

Therefore, applying Lemma 4.1, there exists a $\xi \in (n, n+1)$ such that

$$\begin{aligned}
(n+1)^d - \left(n^{d/(1-p)} + 1 \right)^{1-p} &\geq (n+1)^d - n^d - (1-p)n^{-\frac{dp}{1-p}} \\
&= n^d + d\xi^{d-1} - n^d - (1-p)n^{-\frac{dp}{1-p}} \\
&\geq n^d + dn^{d-1} - n^d - (1-p)n^{-\frac{dp}{1-p}} \\
&= dn^{d-1} - (1-p)n^{-\frac{dp}{1-p}}.
\end{aligned}$$

Consequently, going back to (14),

$$\theta_{f(n)} \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j > \frac{(n^{d/(1-p)} + 1)^{-q}}{1-p} \left(dn^{d-1} - (1-p)n^{-\frac{dp}{1-p}} \right).$$

By Lemma 4.1, there now exists $\xi \in (n^{d/(1-p)}, n^{d/(1-p)} + 1)$ such that

$$\begin{aligned}
\theta_{f(n)} \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j &> \frac{n^{-\frac{dq}{1-p}} - q\xi^{-q-1}}{1-p} \left(dn^{d-1} - (1-p)n^{-\frac{dp}{1-p}} \right) \\
&\geq \frac{n^{-\frac{dq}{1-p}} - qn^{-\frac{d(q+1)}{1-p}}}{1-p} \left(dn^{d-1} - (1-p)n^{-\frac{dp}{1-p}} \right) \\
&\geq \frac{d}{1-p} n^{d-1-\frac{dq}{1-p}} - n^{-\frac{d(p+q)}{1-p}} - \frac{dq}{1-p} n^{d-1-\frac{d(q+1)}{1-p}}.
\end{aligned}$$

Now observe that $d-1-\frac{dq}{1-p} = d(1-\frac{q}{1-p})-1 > \frac{d}{d}-1=0$ and $d-1-\frac{d(q+1)}{1-p} = d(1-\frac{q}{1-p})-\frac{d}{1-p}-1 \leq 3/2-2 = -1/2$. Moreover, $-\frac{d(p+q)}{1-p} < 0$, so the right-hand-side in the equation above is monotone increasing. Therefore,

$$\theta_{f(n)} \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j > \frac{d}{1-p} - 1 - \frac{dq}{1-p} = \frac{1-q}{1-p} \cdot d - 1, \quad \text{for all } n \geq 1.$$

Since $q < p$, we have $1-q > 1-p$. Moreover, $d > 1$. Thus, we may choose $n_0 := 1$ and $\delta := \frac{d(1-q)}{1-p} - 1 > 0$.

We now calculate the modulus φ_2 . By Lemma 4.1, there exists a real number $\xi \in ((n+1)^{d/(1-p)}, (n+1)^{d/(1-p)} + 1)$ such that

$$\begin{aligned}\theta_{f(n)} - \theta_{f(n+1)} &= \left[n^{d/(1-p)} \right]^{-q} - \left[(n+1)^{d/(1-p)} \right]^{-q} \\ &\leq n^{-\frac{dq}{1-p}} - \left((n+1)^{d/(1-p)} + 1 \right)^{-q} \\ &= n^{-\frac{dq}{1-p}} - \left((n+1)^{-\frac{dq}{1-p}} - q \cdot \xi^{-q-1} \right) \\ &\leq n^{-\frac{dq}{1-p}} - (n+1)^{-\frac{dq}{1-p}} + q \cdot \left((n+1)^{d/(1-p)} \right)^{-q-1} \\ &= n^{-\frac{dq}{1-p}} - (n+1)^{-\frac{dq}{1-p}} + q \cdot (n+1)^{-\frac{d(1+q)}{1-p}} \\ &\leq n^{-\frac{dq}{1-p}} - (n+1)^{-\frac{dq}{1-p}} + q \cdot n^{-\frac{d(1+q)}{1-p}}.\end{aligned}$$

Applying once more Lemma 4.1, we see that for some $\xi \in (n, n+1)$,

$$\begin{aligned}\theta_{f(n)} - \theta_{f(n+1)} &= n^{-\frac{dq}{1-p}} - \left(n^{-\frac{dq}{1-p}} - \frac{dq}{1-p} \xi^{-\frac{dq}{1-p}-1} \right) + q \cdot n^{-\frac{d(1+q)}{1-p}} \\ &= \frac{dq}{1-p} \xi^{-\frac{dq}{1-p}-1} + q \cdot n^{-\frac{d(1+q)}{1-p}} \\ &\leq \frac{dq}{1-p} n^{-\frac{dq}{1-p}-1} + q \cdot n^{-\frac{d(1+q)}{1-p}}.\end{aligned}$$

Since $-\frac{d(1+q)}{1-p} = -\frac{dq}{1-p} - \frac{d}{1-p} \leq -\frac{dq}{1-p} - 1$,

$$\theta_{f(n)} - \theta_{f(n+1)} \leq \left(\frac{dq}{1-p} + q \right) n^{-\frac{dq}{1-p}-1} = \frac{q(d+1-p)}{1-p} n^{-\frac{dq}{1-p}-1}. \quad (15)$$

On the other hand,

$$\begin{aligned}\sum_{j=f(n)}^{f(n+1)} \lambda_j &= \sum_{j=f(n)}^{f(n+1)} j^{-p} \leq \int_{f(n)-1}^{f(n+1)-1} j^{-p} dj \\ &= \frac{1}{1-p} \left(\left[(n+1)^{d/(1-p)} - 1 \right]^{1-p} - \left[n^{d/(1-p)} - 1 \right]^{1-p} \right).\end{aligned}$$

Since $1-p > 0$, we conclude

$$\sum_{j=f(n)}^{f(n+1)} \lambda_j \leq \frac{1}{1-p} \left((n+1)^d - \left(n^{d/(1-p)} - 1 \right)^{1-p} \right).$$

By Lemma 4.1, there exists $\xi \in (n, n+1)$ and $\nu \in (n^{d/(1-p)} - 1, n^{d/(1-p)})$ such that for $n \geq 2$

$$\begin{aligned}(1-p) \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j &\leq n^d + d\xi^{d-1} - \left(n^{d/(1-p)} - 1 \right)^{1-p} \\ &\leq n^d + d(n+1)^{d-1} - \left(n^{d/(1-p)} - 1 \right)^{1-p} \\ &= n^d + d(n+1)^{d-1} - n^d + (1-p)\nu^{-p} \\ &\leq n^d + d(n+1)^{d-1} - n^d + (1-p) \left(n^{d/(1-p)} - 1 \right)^{-p} \\ &\leq d(n+1)^{d-1} + (1-p) \cdot 2^p n^{-\frac{dp}{1-p}} \\ &\leq d \cdot 2^{d-1} n^{d-1} + (1-p) \cdot 2^p n^{-\frac{dp}{1-p}}.\end{aligned}$$

Combining this with (15), we get

$$\begin{aligned} (\theta_{f(n)} - \theta_{f(n+1)}) \sum_{j=f(n)}^{f(n+1)} \lambda_j &\leq \left(\frac{q(d+1-p)}{1-p} n^{-\frac{dq}{1-p}-1} \right) \frac{d \cdot 2^{d-1} n^{d-1} + (1-p) \cdot 2^p n^{-\frac{dp}{1-p}}}{1-p} \\ &= \frac{2^{d-1} dq(d+1-p)}{(1-p)^2} n^{d-2-\frac{dq}{1-p}} + \frac{2^p q(d+1-p)}{1-p} n^{-\frac{d(p+q)}{1-p}-1}. \end{aligned}$$

Now observe that $d-2-\frac{dq}{1-p} = d(1-\frac{q}{1-p})-2 \leq \frac{3}{2}(1-\frac{q}{1-p})^{-1}(1-\frac{q}{1-p})-2 = -\frac{1}{2}$. Therefore,

$$\begin{aligned} (\theta_{f(n)} - \theta_{f(n+1)}) \sum_{j=f(n)}^{f(n+1)} \lambda_j &\leq \frac{2^{d-1} dq(d+1-p)}{(1-p)^2} n^{-\frac{1}{2}} + \frac{2^p q(d+1-p)}{1-p} n^{-\frac{d(p+q)}{1-p}-1} \\ &\leq \frac{2^{d-1} dq(d+1-p)}{(1-p)^2} n^{-\frac{1}{2}} + \frac{2^p q(d+1-p)}{(1-p)^2} n^{-\frac{1}{2}} \\ &\leq \frac{2^d dq(d+1-p)}{(1-p)^2} n^{-\frac{1}{2}}. \end{aligned}$$

Therefore, we may take $\varphi_2(\varepsilon) := \left\lceil \left(\frac{2^d dq(d+1-p)}{\varepsilon(1-p)^2} \right)^2 \right\rceil$.

Now, we calculate the modulus φ_3 . To this end, observe that

$$\begin{aligned} \sum_{j=f(n)}^{f(n+1)} \lambda_j^2 &= \sum_{j=f(n)}^{f(n+1)} j^{-2p} \leq \int_{j=f(n)-1}^{f(n+1)-1} j^{-2p} dj \\ &= \begin{cases} \frac{1}{1-2p} \left(\lceil (n+1)^{d/(1-p)} - 1 \rceil^{1-2p} - \lceil n^{d/(1-p)} - 1 \rceil^{1-2p} \right) & , \text{ if } p \neq \frac{1}{2}, \\ \log \lceil (n+1)^{d/(1-p)} - 1 \rceil - \log \lceil n^{d/(1-p)} - 1 \rceil & , \text{ if } p = \frac{1}{2}. \end{cases} \end{aligned}$$

We have to distinguish the cases $p > 1/2$, $p = 1/2$ and $p < 1/2$. For $p > 1/2$, we use the estimate

$$\sum_{j=f(n)}^{f(n+1)} \lambda_j^2 \leq \frac{(n^{d/(1-p)} - 1)^{1-2p}}{2p-1} \leq \varepsilon, \quad \text{for all } n \geq \left(((2p-1)\varepsilon)^{1/(1-2p)} + 1 \right)^{(1-p)/d}.$$

For $p = 1/2$, we see that

$$\sum_{j=f(n)}^{f(n+1)} \lambda_j^2 \leq \log \frac{(n+1)^{d/(1-p)}}{\lceil n^{d/(1-p)} - 1 \rceil} \leq \log \frac{(n+1)^{d/(1-p)}}{n^{d/(1-p)} - 1}.$$

Now observe that $(n+1)^{d/(1-p)} = n^{d/(1-p)} + \frac{d}{1-p} \xi^{\frac{d}{1-p}-1}$ for some $\xi \in (n, n+1)$. Therefore,

$$\begin{aligned} \sum_{j=f(n)}^{f(n+1)} \lambda_j^2 &\leq \log \left(\frac{n^{d/(1-p)} + \frac{d}{1-p} (n+1)^{\frac{d}{1-p}-1}}{n^{d/(1-p)} - 1} \right) \\ &= \log \left(1 + \frac{\frac{d}{1-p} (n+1)^{\frac{d}{1-p}-1} + 1}{n^{d/(1-p)} - 1} \right) \\ &\leq \log \left(1 + 2^{d/(1-p)} \cdot 2^{d/(1-p)} \cdot \frac{\frac{d}{1-p} n^{\frac{d}{1-p}-1}}{n^{d/(1-p)}} \right) \\ &= \log \left(1 + \frac{d \cdot 2^{\frac{2d}{1-p}}}{1-p} \cdot \frac{1}{n} \right) \leq \varepsilon, \quad \text{for all } n \geq \frac{d \cdot 2^{\frac{2d}{1-p}}}{(1-p)(\exp(\varepsilon) - 1)}. \end{aligned}$$

For $p < 1/2$, we see that there exists $\nu \in (n^{d/(1-p)} - 1, n^{d/(1-p)})$ and $\xi \in (n, n+1)$ such that

$$\begin{aligned}
(1-2p) \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j^2 &= \left[(n+1)^{d/(1-p)} - 1 \right]^{1-2p} - \left[n^{d/(1-p)} - 1 \right]^{1-2p} \\
&\leq (n+1)^{\frac{d(1-2p)}{1-p}} - \left(n^{d/(1-p)} - 1 \right)^{1-2p} \\
&= (n+1)^{\frac{d(1-2p)}{1-p}} - n^{\frac{d(1-2p)}{1-p}} + (1-2p)\nu^{-2p} \\
&\leq (n+1)^{\frac{d(1-2p)}{1-p}} - n^{\frac{d(1-2p)}{1-p}} + (1-2p)(n^{d/(1-p)} - 1)^{-2p} \\
&= n^{\frac{d(1-2p)}{1-p}} + \frac{d(1-2p)}{1-p} \xi^{\frac{d(1-2p)}{1-p}-1} - n^{\frac{d(1-2p)}{1-p}} + (1-2p)(n^{d/(1-p)} - 1)^{-2p} \\
&\leq \frac{d(1-2p)}{1-p} n^{\frac{d(1-2p)}{1-p}-1} + (1-2p)(n^{d/(1-p)} - 1)^{-2p}
\end{aligned}$$

Observe that $\frac{d(1-2p)-1+p}{1-p} = \frac{d(1-2p)}{1-p} - 1 < 0$ since $d < \frac{1-p}{1-2p}$. Therefore,

$$\begin{aligned}
\sum_{j=f(n)}^{f(n+1)} \lambda_j^2 &\leq \frac{d}{1-p} n^{\frac{d(1-2p)}{1-p}-1} + (n^{d/(1-p)} - 1)^{-2p} \\
&\leq \varepsilon, \quad \text{for all } n \geq \max \left\{ \left(\frac{2d}{(1-p)\varepsilon} \right)^{\frac{1-p}{d(1-2p)-1+p}}, \left((2/\varepsilon)^{\frac{1}{2p}} + 1 \right)^{(1-p)/d} \right\}.
\end{aligned}$$

Observing that $\theta_n = n^{-q}$ converges to 0 with modulus $\sqrt[q]{1/\varepsilon}$, we summarize the moduli for this choice of the parameter sequences.

1. $n_0 := 1$ and $\delta := \frac{d(1-q)}{1-p} - 1$,
2. $f(n) := \lceil n^{d/(1-p)} \rceil$, where $d := \min \left\{ \frac{1-p}{1-r-p}, \frac{3}{2} \left(1 - \frac{q}{1-p} \right)^{-1} \right\}$,
3. $\varphi_1(\varepsilon) := \sqrt[q]{1/\varepsilon}$,
4. $\varphi_2(\varepsilon) := \left\lceil \left(\frac{2^d d q (d+1-p)}{\varepsilon (1-p)^2} \right)^2 \right\rceil + 1$,
5. $\varphi_3(\varepsilon) := \begin{cases} ((2p-1)\varepsilon)^{\frac{1-p}{d(1-2p)}} + 1 & \text{for } p > \frac{1}{2}, \\ \frac{d \cdot 2^{\frac{2d}{1-p}}}{(1-p)(\exp(\varepsilon)-1)} + 1 & \text{for } p = \frac{1}{2}, \\ \max \left\{ \left(\frac{2d}{(1-p)\varepsilon} \right)^{\frac{1-p}{d(1-2p)-1+p}}, \left((2/\varepsilon)^{\frac{1}{2p}} + 1 \right)^{(1-p)/d} \right\} + 1 & \text{for } p < \frac{1}{2}. \end{cases}$

4.2 Example 2

We begin with the following well-known inequality, whose proof we include for completeness.

Lemma 4.2. For all $x \geq 0$, $\log(1+x) \leq \frac{x}{\sqrt{1+x}}$.

Proof. Define $f : [0, \infty] \rightarrow \mathbb{R}$ by $f(x) := \frac{x}{\sqrt{1+x}} - \log(1+x)$. Then

$$f'(x) = \frac{\sqrt{1+x} - \frac{x}{2\sqrt{1+x}}}{1+x} - \frac{1}{1+x} = \frac{\frac{2+2x-x}{\sqrt{1+x}} - 1}{1+x} = \frac{2+x-\sqrt{1+x}}{(1+x)^{3/2}} \geq 0.$$

Moreover, $f(0) = 0$, so $f(x) \geq 0$ for all $x \geq 0$, whence the claim follows. $\square$

Set $\lambda_n = 1/n$ and $\theta_n = 1/\log \log n$ for $n \geq 3$ and $\lambda_1 = \lambda_2 = \theta_1 = \theta_2 = 0$ (see [1]). Then, we may take $n_0 := 3$, $\delta := 1/2$, $f(n) := n^n$, $\varphi_1(\varepsilon) := \exp \exp(1/\varepsilon)$, $\varphi_2 := \max\{e^4, \exp((1/\varepsilon)^2 - 1) - 1\}$ and $\varphi_3 := \max\{3, \log(2/\varepsilon + 1)\}$. That φ_1 is as required is immediate. Moreover, by Example 1 of [1],

$$\theta_{f(n)} \sum_{j=f(n)}^{f(n+1)} \lambda_j \geq \frac{\log n}{\log n + \log \log n} \geq \frac{\log n}{2 \log n} = \frac{1}{2}, \quad \text{for all } n \geq 3,$$

so n_0 and δ are as required.

Again from [1],

$$1 + \log(n+1) \geq \sum_{j=f(n)}^{f(n+1)} \lambda_j \geq \log n, \quad \text{for all } n \geq 3.$$

Moreover, for $n \geq 3$,

$$\begin{aligned} \log \log(n+1)^{n+1} - \log \log n^n &\leq \log \log \frac{(n+1)^{n+1}}{n^n} = \log \log \left(\left(\frac{n+1}{n} \right)^n (n+1) \right) \\ &\leq \log \log(e(n+1)). \end{aligned}$$

Consequently

$$\begin{aligned} \theta_{f(n)} - \theta_{f(n+1)} &= \frac{1}{\log \log n^n} - \frac{1}{\log \log(n+1)^{n+1}} \\ &= \frac{\log \log(n+1)^{n+1} - \log \log n^n}{(\log \log n^n) \cdot (\log \log(n+1)^{n+1})} \\ &\leq \frac{\log \log(e(n+1))}{(\log \log n^n) \cdot (\log \log(n+1)^{n+1})}. \end{aligned}$$

Now $e^4 \log e^4 = 4e^4 = 3e^4 + e^4 > e^5 + e = e(e^4 + 1)$, so $\log(n \log n) \geq \log(e(n+1))$ for all $n \geq e^4$. Consequently,

$$\begin{aligned} (\theta_{f(n)} - \theta_{f(n+1)}) \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j &\leq \frac{1 + \log(n+1)}{\log(n \log n)} \cdot \frac{\log \log(e(n+1))}{\log((n+1) \log(n+1))} \\ &\leq \frac{\log(e(n+1))}{\log(n \log n)} \cdot \frac{\log(1 + \log(n+1))}{\log(n+1)} \\ &\leq 1 \cdot \frac{\log(1 + \log(n+1))}{\log(n+1)}. \end{aligned}$$

Now, we apply Lemma 4.2

$$\begin{aligned} (\theta_{f(n)} - \theta_{f(n+1)}) \cdot \sum_{j=f(n)}^{f(n+1)} \lambda_j &\leq \frac{\log(n+1)}{\log(n+1) \sqrt{1 + \log(n+1)}} \\ &= \frac{1}{\sqrt{1 + \log(n+1)}} \\ &\leq \varepsilon, \quad \text{for all } n \geq \max\{e^4, \exp((1/\varepsilon)^2 - 1) - 1\}. \end{aligned}$$

Moreover,

$$\begin{aligned} \sum_{j=f(n)}^{f(n+1)} \lambda_j^2 &\leq \int_{j=f(n)-1}^{f(n+1)-1} j^{-2} dj = -2 \left(\left((n+1)^{(n+1)} - 1 \right)^{-1} - (n^n - 1)^{-1} \right) \\ &= \frac{2}{n^n - 1} - \frac{2}{(n+1)^{(n+1)} - 1} \\ &\leq \frac{2}{n^n - 1} \leq \varepsilon, \quad \text{for all } n \geq \max\{3, \log(2/\varepsilon + 1)\}. \end{aligned}$$

References

- [1] R. E. Bruck. A strongly convergent iterative solution of $0 \in U(x)$ for a maximal monotone operator U in Hilbert space. *Journal of Mathematical Analysis and Applications*, 48:114–126, 1974.
- [2] C. Chidume. *Geometric Properties of Banach Spaces and Nonlinear Iterations*. Springer Lecture Notes in Mathematics, 2009.
- [3] C. W. Henson and J. Iovino. Ultraproducts in analysis. In *Analysis and logic (Mons, 1997)*, volume 262 of *London Mathematical Society Lecture Note Series.*, pages 1–110. Cambridge University Press, Cambridge, 2002.
- [4] U. Kohlenbach. *Applied Proof Theory: Proof Interpretations and their Use in Mathematics*. Springer Monographs in Mathematics, 2008.
- [5] U. Kohlenbach. On quantitative versions of theorems due to F.E. Browder and R. Wittmann. *Advances in Mathematics*, 226:2764–2795, 2011.
- [6] D. Körnlein and U. Kohlenbach. Rate of metastability for Bruck’s iteration of pseudocontractive mappings in Hilbert space. *Numerical Functional Analysis and Optimization*, 35(1):20–31, 2014.
- [7] K. Lan and J. Wu. Convergence of approximants for demicontinuous pseudo-contractive maps in Hilbert spaces. *Nonlinear Analysis: Theory, Methods & Applications*, 49(6):737–746, June 2002.
- [8] E. Neumann. Computational problems in metric fixed point theory and their Weihrauch degrees. *Logical Methods in Computer Science*, 11(4):1–44, 2015.